



(elexacaftor/tezacaftor/ivacaftor and ivacaftor)

100 mg/50 mg/75 mg and 150 mg tablets
50 mg/25 mg/37.5 mg and 75 mg tablets
100 mg/50 mg/75 mg and 75 mg oral granules
80 mg/40 mg/60 mg and 59.5 mg oral granules

CONVERSATION STARTERS FOR PEOPLE CONSIDERING TRIKAFTA®



Am I or is my loved one eligible for TRIKAFTA?



Why should I consider TRIKAFTA?



How does TRIKAFTA work?



What are the possible side effects? What kinds of tests are needed before and while taking TRIKAFTA?



What types of improvements might someone experience on TRIKAFTA?



How is TRIKAFTA different from other cystic fibrosis (CF) treatments? How would TRIKAFTA fit into a treatment routine?



When starting TRIKAFTA, should other medicines be adjusted or stopped?

See page 2 for space to write down your notes.

What is TRIKAFTA® (elexacaftor/tezacaftor/ivacaftor and ivacaftor)?

TRIKAFTA is a prescription medicine used for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have at least one copy of the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or another mutation that is responsive to treatment with TRIKAFTA.

Talk to your doctor to learn if you have an indicated CF gene mutation.

It is not known if TRIKAFTA is safe and effective in children under 2 years of age.



IMPORTANT SAFETY INFORMATION

Before taking TRIKAFTA, tell your doctor about all of your medical conditions, including if you:

- are allergic to TRIKAFTA or any ingredients in TRIKAFTA. See the Patient Information for a list of ingredients
- have kidney problems
- have or have had liver problems
- are pregnant or plan to become pregnant. It is not known if TRIKAFTA will harm your unborn baby. You and your doctor should decide if you will take TRIKAFTA while you are pregnant
- are breastfeeding or planning to breastfeed. It is not known if TRIKAFTA passes into your breast milk. You and your doctor should decide if you will take TRIKAFTA while you are breastfeeding

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#), including [Patient Information](#).



(elexacaftor/tezacaftor/ivacaftor and ivacaftor)

100 mg/50 mg/75 mg and 150 mg tablets
50 mg/25 mg/37.5 mg and 75 mg tablets
100 mg/50 mg/75 mg and 75 mg oral granules
80 mg/40 mg/60 mg and 59.5 mg oral granules

YOUR NOTES



Use this space to write down any notes or additional questions you have for your healthcare provider.

Lined writing area for notes.



(eleacaftor/tezacaftor/ivacaftor
and ivacaftor)

100 mg/50 mg/75 mg and 150 mg tablets
50 mg/25 mg/37.5 mg and 75 mg tablets
100 mg/50 mg/75 mg and 75 mg oral granules
80 mg/40 mg/60 mg and 59.5 mg oral granules

IMPORTANT SAFETY INFORMATION (Continued)



Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

TRIKAFTA may affect the way other medicines work, and other medicines may affect how TRIKAFTA works. The dose of TRIKAFTA may need to be adjusted when taken with certain medicines. Ask your doctor or pharmacist for a list of these medicines if you are not sure.



Especially tell your doctor if you take:

- antibiotics such as rifampin (RIFAMATE[®], RIFATER[®]) or rifabutin (MYCOBUTIN[®])
- seizure medicines such as phenobarbital, carbamazepine (TEGRETOL[®], CARBATROL[®], EQUETRO[®]), or phenytoin (DILANTIN[®], PHENYTEK[®])
- St. John's wort
- antifungal medicines including ketoconazole, itraconazole (such as SPORANOX[®]), posaconazole (such as NOXAFIL[®]), voriconazole (such as VFEND[®]), or fluconazole (such as DIFLUCAN[®])
- antibiotics including telithromycin, clarithromycin (such as BIAXIN[®]), or erythromycin (such as ERY-TAB[®])



What should I avoid while taking TRIKAFTA?

- Avoid food or drink that contains grapefruit while you are taking TRIKAFTA



What are the possible side effects of TRIKAFTA?

TRIKAFTA can cause serious side effects, including:



Liver damage and worsening of liver function in people with severe liver disease that can be serious and may require transplantation. Liver damage has also happened in people without liver disease



High liver enzymes in the blood, which is a common side effect in people treated with TRIKAFTA. These can be serious and may be a sign of liver injury. Your doctor will do blood tests to check your liver:

- before you start TRIKAFTA
- every 3 months during your first year of taking TRIKAFTA
- every year while you are taking TRIKAFTA



(elexacaftor/tezacaftor/ivacaftor and ivacaftor)

100 mg/50 mg/75 mg and 150 mg tablets
50 mg/25 mg/37.5 mg and 75 mg tablets
100 mg/50 mg/75 mg and 75 mg oral granules
80 mg/40 mg/60 mg and 59.5 mg oral granules

IMPORTANT SAFETY INFORMATION (Continued)



What are the possible side effects of TRIKAFTA (continued)?

Your doctor may do blood tests to check the liver more often if you have had high liver enzymes in your blood in the past.

Call your doctor right away if you have any of the following symptoms of liver problems:

- pain or discomfort in the upper right stomach (abdominal) area
- yellowing of your skin or the white part of your eyes
- loss of appetite
- nausea or vomiting
- dark, amber-colored urine



Serious allergic reactions have happened to people who are treated with TRIKAFTA. Call your healthcare provider or go to the emergency room right away if you have any symptoms of an allergic reaction. Symptoms of an allergic reaction may include:

- rash or hives
- tightness of the chest or throat or difficulty breathing
- swelling of the face, lips, and/or tongue, or difficulty swallowing
- light-headedness or dizziness



Abnormality of the eye lens (cataract), which has happened in some children and adolescents treated with TRIKAFTA. If you are a child or adolescent, your doctor should perform eye examinations before and during treatment with TRIKAFTA to look for cataracts



The most common side effects of TRIKAFTA include:

- headache
- upper respiratory tract infection (common cold) including stuffy and runny nose
- stomach (abdominal) pain
- diarrhea
- rash
- increase in liver enzymes
- increase in a certain blood enzyme called creatine phosphokinase
- flu (influenza)
- inflamed sinuses
- increase in blood bilirubin

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of TRIKAFTA. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#), including [Patient Information](#).



TRIKAFTA is manufactured for Vertex Pharmaceuticals Incorporated. TRIKAFTA, the TRIKAFTA logo, Vertex, and the Vertex triangle logo are registered trademarks of Vertex Pharmaceuticals Incorporated. All other trademarks referenced herein are the properties of their respective owners. © 2023 Vertex Pharmaceuticals Incorporated | VXR-US-28-1900042 (v6.0) | 09/2023